

POSTER PRESENTATION

Open Access


Neurodynamic regimes of phase relation and behavior in robotic models

Bruno A Santos^{1,3*}, Xabier E Barandiaran², Phil Husbands¹

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

Both oscillatory phenomena and the dynamics of integration and segregation in the brain are fields of intense study in current computational neuroscience. It is often suggested that moments of phase-locking (as moments of integration) between functionally distinct neuronal groups represent meaningful functional brain states; and moments of phase-scattering (as moments of segregation) represent transitions between such states. The main evidences for such claim come from investigations on the well-known visual binding problem [1]. Despite the evidences supporting a functionally privileged status for phase locking integration (as a neural signature of cognitive brain activity) an adequate balance between integration and segregation has also been considered essential [2]. We explore how, rather than mere phase-locking, the global *regime of phase relation* might become necessary to generate functional behaviour, i.e.

the whole phase dynamic which might include moments of integration (by phase-locking) and segregation (by phase scattering). Our proposal to investigate the whole phase relation dynamics is motivated by the phenomenon of *relative coordination* observed in the movements of fish fins described by von Holst [3] and latter developments of Haken and Kelso's approach to coordination dynamics [4]. In dynamical system terms, by "relative coordination" von Holst means the situation when the phase relation among components is constantly moving in a transient dynamics where some regions of the phase space have low potential energy (moments of phase-locking) and others high potential energy (moments of phase-scattering). Using a genetic algorithm we optimized the parameter of two robotic models performing phototaxis. In one model the agent is controlled by the extended-HKB equation [4] (Fig. 1-



* Correspondence: b.santos@sussex.ac.uk

¹CCNR - University of Sussex, Brighton UK

Full list of author information is available at the end of the article

A-,1B) and in the other one by a network of five coupled Kuramoto oscillators [5] (Fig. 1-C,1D). Both types of controllers were chosen as paradigmatic of oscillatory modeling in computational neuroscience.

Due to the simplicity of the model we were able to analyze the whole phase dynamic and found different dynamical regimes of phase relation (expressed in terms of phase density) and their behavioral correlates. Despite the strong simplifying assumptions involved, the theoretical results shown here suggest a “different/new” approach to analyze empirical data in neuroscience where regimes of phase relation might be as relevant as phase locking events for how brain-environment dynamics give rise to behaviour. We conclude with some expansions of phase density and phase coherence measurements for bigger oscillatory networks in sensorimotor systems.

Author details

¹CCNR - University of Sussex, Brighton UK. ²CREA - Polytechnique/CNRS, Paris, France. ³LSI, CEFET-MG, Belo Horizonte, Brazil.

Published: 18 July 2011

References

1. Singer W: **Visual feature integration and the temporal correlation hypothesis.** *Annu. Rev. Neurosci* 1995, **18**:555-586.
2. Tononi G, Sporns O, Edelman GM: **A measure for brain complexity: relating functional segregation and integration in the nervous system.** *Proceedings of the National Academy of Sciences of the United States of America* 1994, **91**(11):5033-5037.
3. von Holst E: **Die relative Koordination als Phaenomen und als Methode zentralnervoerer Funktionsanalyse.** *Ergebnisse Physiol* 1939, **42**:228-306.
4. Haken H, Kelso JAS, Bunz H: **A theoretical model of phase transitions in human hand movements.** *Biol Cybern* 1985, **51**(5):347-356.
5. Kuramoto Y: **Chemical oscillations, waves, and turbulence.** Springer, New York; 1984.

doi:10.1186/1471-2202-12-S1-P155

Cite this article as: Santos et al.: Neurodynamic regimes of phase relation and behavior in robotic models. *BMC Neuroscience* 2011 **12**(Suppl 1):P155.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

